



## Phytochemical Constituents of *Pleurotus ostreatus*

Hasina Begum Bina<sup>1</sup>, SM Mizanur Rahman<sup>2</sup>, Anath Chandra Roy<sup>3</sup>, Tofail Ahmad Chowdhury<sup>4</sup>, Md. Hasanur Rahman<sup>5</sup>, Abdullah Al-Mansur<sup>6</sup>, Md. Eyazul Haque<sup>7\*</sup>

<sup>1-5</sup> Department of Chemistry, University of Dhaka, Dhaka-1000, Bangladesh

<sup>6</sup> Institute of National Analytical Research and Service, BCSIR, Dhaka, Bangladesh

<sup>7</sup> Department of Chemistry, Bangladesh University of Engineering and Technology, Dhaka, Bangladesh

DOI: <https://doi.org/10.33545/26646781.2019.v1.i2a.10>

### Abstract

The chromatographic separation of the dichloromethane extract of *Pleurotus ostreatus*, fatty acid derivatives like trans-4-acetoxy-5-hydroxy-5-pentadecanoyl-2-cyclopenten-1-one and trans-4-acetoxy-5-hydroxy-5-(1-acetoxypentadecyl)-2-cyclopenten-1-one were isolated. The structures of these compounds have been determined by <sup>1</sup>H, <sup>13</sup>C, DEPT-135 NMR and GC-MS spectral data.

**Keywords:** *Pleurotus ostreatus*, Fatty acid, NMR, GC-MS

### 1. Introduction

Mushrooms are one type of fungi. In Bangladesh it grows wild all over the country under favorable climatic conditions. Most of the mushrooms have nutritional and medicinal <sup>[1,2,3,4]</sup> value. FAO has recommended <sup>[5]</sup> edible mushroom as a source of protein nutrition of developing countries where malnutrition is a serious problem. Mushrooms contain 19-35% proteins <sup>[6]</sup> on dry weight basis as compared to 7.3% in rice, 13.2% in wheat, and 25.2% in milk. Mushroom contains all the nine essential amino acids to a considerable amount. Oyster mushroom (*Pleurotus ostreatus*) belongs to the *Pleurotaceae* family, is an edible mushroom used as nutritious and delicious food. Now-days it is cultivated in many places of the country. It was reported <sup>[6]</sup> that *Pleurotus ostreatus* rich in protein and as well as medicinal importance. In this study, is described the isolation and structure elucidation of two fatty acid derivatives.

### 2. Materials and methods

#### 2.1 General Experimental Section

Freshly distilled solvents were used for extraction, isolation and purification. Evaporations were performed under reduced pressure on a Buchii rotary evaporator. Melting point was determined on an electrochemical micro-melting point apparatus. The UV, IR (KBr) spectra were recorded on a Shimadzu UV-168A and Shimadzu IR-470A spectrophotometer, respectively. The <sup>1</sup>H- and <sup>13</sup>C-NMR spectra were taken in deuterated methanol (CD<sub>3</sub>OD) with TMS as an internal standard by using a high-resolution 400 MHz spectrometer.

#### 2.2 Plant Material

The fruit body of mushroom "*Pleurotus ostreatus*" has been collected from the National Mushroom Development and Extend Center, Savar of Dhaka district and these were identified from the department of Botany, Dhaka University.

#### 2.3 Extraction and Isolation of compound 1 & 2

The fruit body of mushroom was dried under mild sunlight and then at 400 C in an oven. Afterwards the mushrooms were

powdered in a grinding machine (~ 200 mesh). The powdered Mushroom sample (913g) was successively and exhaustively extracted in a soxhlet apparatus with *n*-hexane followed by dichloromethane. The dichloromethane extract was concentrated (0.67g) and subjected to vacuum liquid chromatography (VLC) over TLC grade silica gel (60, P<sub>254</sub>). The column was eluted initially with *n*-hexane followed by the mixture of *n*-hexane and dichloromethane of increasing amount of dichloromethane and finally with methanol. These elutes were collected in a series of test tubes with 25 ml in each fraction. Based on the similar behavior after monitoring each fraction by TLC these elutes were combined to yield eight different fractions as F<sub>1</sub>, F<sub>2</sub>, F<sub>3</sub>, F<sub>4</sub>, F<sub>5</sub>, F<sub>6</sub>, F<sub>7</sub> and F<sub>8</sub>. The fractions (F<sub>1</sub>-F<sub>8</sub>) were concentrated and allowed to stand at room temperature for several days. Precipitated was obtained from the fraction F<sub>6</sub> and it was washed with petroleum ether followed by chloroform and dichloromethane successively. After dissolving the solid particles in methanol that was applied in TLC micro glass slide under different solvent systems and viewed in an iodine chamber. In a particular solvent system where the ratio of dichloromethane and methanol was 9:1 two bright prominent spot was detected. For further purification the solid was applied to column over column grade silica gel (kiesel gel 60, mesh 70-230). The column was eluted with mixture of dichloromethane and methanol by gradual increasing the polarity of the mixture. The eluents were collected in an amount of about 5 ml in a series of test tubes. Based on the similar behavior after monitoring each fraction by TLC these elutes were combined to yield six different fractions. All these fractions (B<sub>1</sub>-B<sub>6</sub>) were concentrated and allowed to stand at room temperature for several days. A reddish solid was obtained from the sub-fractions B<sub>3</sub> (75.2 mg) and B<sub>6</sub> ( 72.5 mg). These crystals were washed separately with the solvent dichloromethane and the solutions were kept at room temperature in two separate vials for recrystallization. Reddish solid precipitated were obtained in both the cases and these compounds were designated as B<sub>3</sub> and B<sub>6</sub>, respectively. After dissolving the crystals in dichloromethane in

each cases that was applied in TLC micro glass slide in the solvent systems dichloromethane: methanol where the ratio is 95 : 05. In both the cases a single spot was detected when viewed in an iodine chamber. The spot of B<sub>3</sub> as the compound 1 appears as violet color when vanillin-sulphuric acid reagent was spraying on the applied TLC plate followed by heating at 110°C in an electric oven with the R<sub>f</sub> value of 0.66 where in case of B<sub>6</sub> as compound 2 the color of the spot was appeared as black under same condition with R<sub>f</sub> value of 0.45.

**Compound 1:** Reddish solid, R<sub>f</sub> value of 0.66; IR (KBr)  $\lambda$  cm<sup>-1</sup>: 3500-3600, 2987, 1710, 1637, 1421, 1363, and 720; <sup>1</sup>H-NMR (400 Mz, CDCl<sub>3</sub>): 7.42 (3-H), 7.25 (2-H), 5.69 (4-H), 4.06-4.14 (5-OH), 2.73 (7a-H), 2.308 (7b-H), 2.034 (4-OAc), 1.620 (8-H), 1.24-1.32 (9-H-19-H), and 0.876 (20-H); <sup>13</sup>C-NMR (400 Mz, CDCl<sub>3</sub>): 14.05 (CH<sub>3</sub>-C-20), 22.06 (C-19), 22.54 (C-18), 22.68 (C-17), 22.67 (C-16), 25.47 (C-15), 27.39 (C-14), 28.87 (C-13), 29.04 (C-12), 29.28 (C-11), 29.69 (C-10), 31.51 (C-9), 31.70 (C-8), 124.54 (C-2), 133.82 (C-3), 21.11 (CH<sub>3</sub> at C-4), 33.57, (CH<sub>2</sub> at C-7), 73.66 (CH at C-4), and 77.35 (quaternary carbon C-5); EIMS: M<sup>+</sup> m/z (Rel. int.): [M+1] 415 (12.5), 327.2 (27.5), 281.3 (31.1), 241.2 (100), 2.37.2 (38.7), 197.3 (33.7), 143.2 (27.5) and 71.1 (46.2).


**Compound 2:** Reddish solid; R<sub>f</sub> value of 0.45; IR (KBr)  $\lambda$  cm<sup>-1</sup>: 3400, 2929, 1710, 1637, 1425, 1359, and 720; <sup>1</sup>H-NMR (400 Mz, CDCl<sub>3</sub>): 7.25 (3-H), 5.58 (2-H) and or 4-H), 5.17 (6-H), 3.31 (5-OH), 2.16 (4-OAc), 2.03 (6-OAc), 1.62 (7-H), 1.20-1.46 (8-H and 9-H), and 0.87 (20-H); <sup>13</sup>C-NMR (400 Mz, CDCl<sub>3</sub>): 14.04 (CH<sub>3</sub>-C-20), 22.56 (C-19), 24.68 (C-18), 27.40 (C-17), 28.91 (C-16 and C-7), 29.08 (C-15), 29.31(C-14), 29.51(C-13), 29.72 (C-12), 30.91 (C-11), 31.53 (C-10), 31.73 (C-9), 33.60 (C-8), 124.57 (C-2), 133.81 (>C=C at C-3), 22.56 (acetoxyl CH<sub>3</sub> at C-4), 76.71 (CH at C-6), 77.03 (CH at C-4) and 206.96 (>C=O at C-1); M<sup>+</sup> m/z (Rel. int.): [M+1] 270 (27.2), 227.2 (40.0), 143.2 (47.1), 88.2 (85.2), 74.1 (100), 55.2 (60.1), 43.1 (72.5) and 41.1 (46.3).

### 3. Results & Discussion

The thin layer chromatographic examination of the isolated compound 1 from the dichloromethane extract of the *Pleurotus ostreatus* followed by successive chromatographic separation like VLC and adsorption column chromatography showed as violet color when vanillin-sulphuric acid reagent was spraying on the applied TLC plate followed by heating at 110°C in an electric oven with the R<sub>f</sub> value of 0.66. The compound 1 was obtained as reddish solid. Its IR spectrum showed <sup>[7]</sup> an absorption peak at (3600-3500) cm<sup>-1</sup> was assignable to a hydroxyl group (-OH) whereas absorption at 2987, and 1363 cm<sup>-1</sup> were assigned due to the presence of aliphatic C-H stretching of -CH<sub>3</sub> and >CH- groups, respectively. A single absorption band at 1637 cm<sup>-1</sup> was demonstrated for C=C groups. The absorption peaks at 1421 cm<sup>-1</sup> was suggestive for -CH<sub>2</sub>- group. The absorption band at 1710 cm<sup>-1</sup> was suggestive for >C=O stretching of the carbonyl group. The other absorption peaks at 720 cm<sup>-1</sup> was the indicative of the long chain of hydrocarbons. The proton NMR study of the isolated compound 1 showed <sup>[8]</sup> a downfield doublet of doublet at  $\delta$  7.420 and 7.250 ppm which was suggestive of the presence of olefinic proton at (C-2 and C-3), respectively. The doublet-doublet at  $\delta$  5.69 ppm was assigned for the proton at C-4. A broad band multiple at  $\delta$  4.06-4.14 ppm was suggestive of an oxymethine proton at C-5. The singlet at  $\delta$  2.034 ppm was assigned for the proton of acetoxy group at C-4. The multiplet at

$\delta$  1.62 and triplet at  $\delta$  0.876 ppm were assigned for the methylene proton at C-8 and methyl proton at C-20, respectively. The chemical shift at  $\delta$  1.245 to 1.320 ppm was appearance as multiplet for the methylene proton that linked from C-9 and C-19 positions. The resonance at  $\delta$  2.73 and 2.308 ppm were assigned for the protons linked to C-7A and C-7B, respectively. The <sup>13</sup>C-NMR spectrum of the isolated compound 1 revealed <sup>[8, 9]</sup> the presence of 22 carbons. The chemical shift at  $\delta$  14.05 ppm was assigned to a methyl carbon at position C-20. The signals at  $\delta$  22.06, 22.54, 22.68, 24.67, 25.47, 27.39, 28.87, 29.04, 29.28, 29.69, 31.51, 31.70 and at 33.57 ppm were assigned for the methylene carbons at the positions 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8 and 7, respectively. The signal for the chemical shifts at  $\delta$  124.54 (C-2) and 133.82 ppm (C-3) were assigned for the C=C between the carbons at position 2 and 3. The resonance at  $\delta$  21.11 ppm was assigned for acetoxy methyl carbon attached to C-4. The resonance at  $\delta$  73.66 ppm was appropriate for the secondary carbon at position 4 while the chemical shift at  $\delta$  77.35 ppm was assigned for the quaternary carbon at position 5. The mass spectroscopic analysis of the compound 1 assigned the molecular ion of the compound m/z is 415, however other peaks could not be explained. The thin layer chromatographic examination of the isolated compound 2 from the dichloromethane extract of the oyster mushroom followed by successive chromatographic separation like VLC and adsorption column chromatography showed as black color when vanillin-sulphuric acid reagent was spraying on the applied TLC plate followed by heating at 110°C in an electric oven with the R<sub>f</sub> value of 0.45. The compound was obtained as reddish solid. Its IR spectrum showed <sup>[7]</sup> an absorption band at 3400 cm<sup>-1</sup> was assignable to a hydroxyl group (-OH) whereas absorption band at 2929, and 1359cm<sup>-1</sup> were assigned due to the presence of aliphatic C-H stretching of -CH<sub>3</sub> and >CH- groups, respectively. A single absorption band at 1637 cm<sup>-1</sup> was demonstrated for the presence of C=C groups. The absorption peaks at 1425 cm<sup>-1</sup> was suggestive for -CH<sub>2</sub>- group. The absorption band at 1710 cm<sup>-1</sup> was suggestive for >C=O stretching of the carbonyl group. The other absorption peaks at 720 cm<sup>-1</sup> was the indicative of the long chain of hydrocarbons. The proton NMR study of the isolated compound 2 showed <sup>[8]</sup> a downfield chemical chemical shift at  $\delta$  5.58 and 7.25 ppm which were suggestive <sup>[9]</sup> of the presence of olefinic proton at (C-2 and C-3), respectively. The chemical shift displayed at  $\delta$  5.17 ppm was assigned for the proton at C-6. A resonance at  $\delta$  3.31 ppm was suggestive of an oxymethine proton at C-5. The resonances at  $\delta$  2.16 and 2.03 ppm was assigned for the proton of the acetoxy groups that linked at C-4 and C-6, respectively. The multiplet at  $\delta$  1.62 and triplet at  $\delta$  0.87 ppm were assigned for the methylene proton at C-7 and methyl proton at C-20, respectively. The chemical shift at  $\delta$  1.20 to 1.47 ppm was appearance as multiplet for the methylene protons that linked from C-8 to C-19 positions. The <sup>13</sup>C-NMR spectrum of the isolated compound 2 revealed <sup>[8, 9]</sup> the presence of 24 carbons. The chemical shift at  $\delta$  14.04 ppm was assigned to a methyl carbon at position C-20. The signals at  $\delta$  22.56, 24.68, 27.40, 28.91, 29.08, 29.31, 29.51, 29.72, 30.91, 33.53, 31.73, 33.60 and at 28.91 ppm were assigned for the methylene carbons at the positions 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8 and 7, respectively. Resonances at  $\delta$  124.57 and at 133.81 ppm were assigned for the olefinic carbons at position 2 and 3, respectively. The resonance at  $\delta$  22.57 ppm was assigned for acetoxy methyl

carbon attached to C-4. The resonances at  $\delta$  76.72 and at 77.03 ppm were appropriate for the secondary carbon at position 6 and 4, respectively, while the chemical shift at  $\delta$  77.35 ppm was assigned for the quaternary carbon at position 5. The chemical shift at  $\delta$  206.96 ppm was suggestive for the presence of  $>C=O$  linked to C-1. The mass spectroscopic analysis of the compound 2 assigned the molecular ion of the compound  $m/z$  is 270.



**Fig 1**

#### 4. Conclusions

A phytochemical analysis has been carried out on the fruit body of mushroom. During this investigation two compounds were isolated from this fungus. All the compounds were identified preliminarily by chemical methods and then the structural elucidation of the compounds were performed by various spectroscopic methods like IR, <sup>1</sup>H-NMR, <sup>13</sup>C-NMR, DEPT-135 and GC-MS and were confirmed by published literatures. These data were identified as compound 1 and 2 are trans-4-acetoxy-5-hydroxy-5-pentadecanoyl-2-cyclopenten-1-one and trans-4-acetoxy-5-hydroxy-5-(1-acetoxypentadecyl)-2-cyclopenten-1-one respectively.

#### 5. Acknowledgments

The authors are grateful to the Department of Chemistry, University of Dhaka and Atomic Energy Centre, Dhaka, Bangladesh for their financial support.

#### 6. References

1. Marmion VJ et.al. The Death of Claudius, the Royal Society of Medicine, 2002.
2. Periasamy K. Novel Antibacterial Compounds Obtained from Some Edible Mushrooms, *Int. J. Med. Mush.*, 2005; **7**: 443-444.
3. Bobeq P, Galbavy S. Hypocholesteremic and Antiatherogenic Effect of Oyster Mushroom (*Pleurotus ostreatus*) in Rabbit, *Nahrung*, 1999; **43**: 339-342.
4. Wasser SP. Medicinal Mushrooms as a Source of Antitumor and Immunomodulating Polysaccharides, *Appl. Microbial. Biotechnol.*, 2002; **60**: 258-274.
5. Mushrooms by Country. <http://www.fao.org>. 7 September, 2008.
6. Chang SP, Miles G. Mushrooms, Cultivation, Natural Value, Medicinal Effect and Environmental Impact, CRC Press, 2004.
7. Lambert, *et al.* Spectroscopy of Organic Chemistry, 1987.
8. Pavia DL, Lampman GL, and Kriz GS. Introduction to Spectroscopy, 3rd edition, USA, 2001.
9. World Intellectual Property Organization. <https://www.wipo.int>. 10 August, 2018.