


Scheme 1: Reaction scheme preparation of tris (2-((E)-1-hydrazonoethyl)-4,5-dimethyl phenoxy)iron.

Preparation of Fe complex with (E)-4,5-dimethyl-2-(1-(2-phenylhydrazono)ethyl)phenol metal complex Or tris(4,5-dimethyl-2-((E)-1-(2-phenylhydrazono)ethyl)phenoxy)iron:

(E)-4,5-dimethyl-2-(1-(2-phenylhydrazono)ethyl)phenol were synthesized following published procedure¹⁷. The transition metal complex tris(4,5-dimethyl-2-((E)-1-(2-phenylhydrazono)ethyl)phenoxy)iron was prepared by dissolution of (E)-4,5-dimethyl-2-(1-(2-phenylhydrazono)ethyl)phenol ligand (0.0103M) in ethanol at hot condition, an ethanolic solution (0.0034M) of the Ferrous Sulfate was added drop wise with

constant stirring and refluxed for 4 hrs. The resulting reaction mixture was cooled to room temperature and maintains pH 7 to 8 by addition of the 0.10 N sodium hydroxide. Thus blue color precipitate was formed. The resultant product was filtered, repeatedly washed with ethanol and dried at 50°C to 55°C temperature. The compound was re-crystallized in ethanol for purification and improving description. Reaction scheme preparation of tris (4,5-dimethyl-2-((E)-1-(2-phenylhydrazono)ethyl)phenoxy) iron is given below Scheme 2.




Scheme 2: Reaction scheme preparation of tris(4,5-dimethyl-2-((E)-1-(2-phenyl hydrazono)ethyl)phenoxy)iron

Preparation of Fe complex with (E)-2-(1-(2-(2,4-dinitrophenyl) hydrazono) ethyl)-4,5-dimethylphenol metal complex Or tris(2-((E)-1-(2-(2,4-dinitrophenyl) hydrazono) ethyl)-4, 5-dimethylphenoxy)iron

(E)-2-(1-(2-(2,4-dinitrophenyl)hydrazono)ethyl)-4,5-dimethylphenol were synthesized following published procedure [18]. The transition metal complex tris(2-((E)-1-(2-(2,4-dinitrophenyl) hydrazono)ethyl)-4,5-dimethylphenoxy)iron was prepared by dissolution of (E)-2-(1-(2-(2,4-dinitrophenyl)hydrazono)ethyl)-4,5-dimethylphenol ligand (0.0087M) in ethanol at hot condition, an ethanolic solution

(0.0028M) of the metal salt Ferrous Sulfate was added drop wise with constant stirring and refluxed for 5 hrs. The resulting reaction mixture was cooled to room temperature and maintains pH 7 to 8 by addition of the 0.10 N sodium hydroxide. Thus brown color precipitate was formed. The resultant product was filtered, repeatedly washed with ethanol and dried at 60°C to 65°C temperature. The compound was re-crystallized in ethanol for purification and improving description. Reaction scheme preparation of tris (2-((E)-1-(2-(2,4-dinitrophenyl) hydrazono) ethyl)-4,5-dimethylphenoxy)iron is given below Scheme 3.



Scheme 3: Reaction scheme preparation of tris(2-((E)-1-(2-(2,4-dinitrophenyl) hydrazono)ethyl)-4,5-dimethylphenoxy)iron.

Results and Discussion

The Fe-Metal complexes were analyzed by Infrared spectroscopy, Ultraviolet-visible spectroscopy, ^1H NMR, Carbon-Hydrogen-Nitrogen-Sulphur elemental analyzer, Decomposition temperature/Melting point, Molar Conductance,

Magnetic susceptibility and Iron content by Gravimetric technique. Decomposition temperature / Melting points were recorded using Veego Scientific Device (Model: VMP-AD) in open capillaries and were uncorrected. Some physical properties of Fe-Metal complexes are mentioned in Table 1.

Table 1: Physical properties

Ni-Metal Complex	Chemical Formula and Molecular formula	Color	Solubility	Decomposition Temperature	Yield
tris(2-((E)-1-hydrazonoethyl)-4,5 dimethylphenoxy)iron	$\text{C}_{30}\text{H}_{39}\text{N}_6\text{FeO}_3$ and 587.51	green	- Soluble in methanol, ethanol, Acetonitrile DMSO, DMF - Insoluble in water	> 300 °C	89.67 %
tris(4,5-dimethyl-2-((E)-1-(2-phenyl hydrazono)ethyl)phenoxy)iron	$\text{C}_{48}\text{H}_{51}\text{N}_6\text{FeO}_3$ and 815.80	blue	- Soluble in ethanol, methanol, DMSO, DMF - Insoluble in water	> 300°C	92.00 %
tris(2-((E)-1-(2-(2,4 dinitrophenyl)hydrazono)ethyl)-4,5 dimethylphenoxy)iron	$\text{C}_{48}\text{H}_{45}\text{N}_{12}\text{FeO}_{15}$ and 1085.79	brown	- Soluble in ethanol, methanol, DMSO, DMF - Insoluble in water	> 300°C	76.63%


Infrared analysis

The IR spectra of Fe-Metal complexes were recorded in the region of 4000–400 cm^{-1} using FTIR spectrometer of model Agilent Resolutions Pro by direct sampling method.

The IR spectral data along with the possible assignments of Fe-Metal complexes are provided in Table 2, followed by IR spectra performed by IR direct solid method, use of Agilent Resolutions Pro and IR spectrum are showed in Figure 1, 2, 3.

Table 2: Hydrazone Iron metal complex IR frequency, cm^{-1}

Bond/functional group	frequency, cm^{-1}		
	Tris (2-((E)-1-hydrazonoethyl)-4,5-dimethyl phenoxy) iron	Tris (4,5-dimethyl-2-((E)-1-(2-phenyl hydrazono)ethyl) phenoxy) iron	Tris (2-((E)-1-(2-(2,4-dinitrophenyl) hydrazono)ethyl)-4,5-dimethyl phenoxy)iron
N-H	3218	3285	3296
C-H	3032	3098	3045
C-H stretch aromatics	2971	2918	2965
C=N	1612	1618	1614
C-C	1548	1506	1507
C=C (aromatic ring)	1494	1493	1418
N-O	---	---	1331
C-O	1257	1248	1261
N-N	1194	1142	1115
O-Fe	Below 600	Below 600	Below 600

**Fig 1:** IR of tris(2-((E)-1-hydrazonoethyl)-4,5-dimethylphenoxy)iron




Fig 2: IR of tris (4,5-dimethyl-2-((E)-1-(2-phenylhydrazono)ethyl) phenoxy)iron




Fig 3: IR of tris (2-((E)-1-(2-(2,4-dinitrophenyl)hydrazono)ethyl)-4,5-dimethylphenoxy)iron



UV analysis

The UV spectra of the Fe-Metal complexes in methanol were recorded on SPECORD 50 PLUS-233 H 1409C Spectro photo meter, using a quartz cell of 1 cm optical path where methanol

was used as a blank. The UV spectrum is shown in Figure 4, 5, 6. The spectra shows λ (bands maximum in nm) are provided in Table 3.

Table 3: Ni-Metal Complex

	Wavelength nm
tris(2-((E)-1-hydrazonoethyl)-4,5 dimethylphenoxy)iron	254, 312,790,850
tris(4,5-dimethyl-2-((E)-1-(2 phenylhydrazono) ethyl) phenoxy) iron	281,354,718,854
tris(2-((E)-1-(2-(2,4-dinitrophenyl)hydrazono)ethyl)-4,5-dimethyl phenoxy)iron	245, 315,728,856

**Fig 4:** UV-Visible Spectra of tris(2-((E)-1-hydrazonoethyl)-4,5-dimethylphenoxy)iron**Fig 5:** UV-Visible Spectra of tris(4,5-dimethyl-2-((E)-1-(2-phenylhydrazono)ethyl)phenoxy)iron




Fig 6: UV-Visible Spectra of tris(2-((E)-1-(2-(2,4-dinitrophenyl)hydrazono)ethyl)-4,5-dimethylphenoxy)iron

Elemental analysis (CHNS)

Formation of Fe-Metal complexes were further confirmed by elemental analysis which was recorded by Vario MICRO CHNS analyzer. The elemental analysis data of Cu-Metal complexes and Cu is confirmed by Gravimetric technique is summarized in Table 4.

Magnetic susceptibility

The magnetic moment (μ_{eff}) values at room temperature for Fe (III) complexes complexes was in the range of 4.5 to 5.3 B.M. These values are expected for Octahedral geometry of Fe (III) complexes at room temperature and further supported by electronic spectral data is summarized in Table 5.

Table 5: Magnetic susceptibility

Fe-Metal complex	μ_{eff} value
tris(2-((E)-1-hydrazoneethyl)-4,5-dimethylphenoxy)iron	4.77
tris(4,5-dimethyl-2-((E)-1-(2-phenylhydrazono) ethyl) phenoxy) iron	4.98
tris(2-((E)-1-(2-(2,4-dinitrophenyl)hydrazono)ethyl)-4,5-dimethyl phenoxy)iron	5.06

Molar Conductance ($\text{mho.cm}^2.\text{mol}^{-1}$)

The molar conductance values at room temperature were measured in Methanol (10^{-3} M) on an Elico digital direct reading conductivity meter model CM-180 is summarized in Table 6.

Table 6: Molar Conductance

Fe-Metal complex	Molar Conductance $\text{mhos cm}^2 \text{mol}^{-1}$
tris(2-((E)-1-hydrazoneethyl)-4,5 dimethyl phenoxy) iron	0.090
tris(4,5-dimethyl-2-((E)-1-(2-phenylhydrazono)ethyl) phenoxy)iron	0.099
tris(2-((E)-1-(2-(2,4-dinitrophenyl) hydrazono)ethyl)-4,5-dimethyl phenoxy)iron	0.097

3.6 ^1H NMR

We tried all Fe-complex ^1H NMR and observe the spectrum of the complex differs from that of the free ligand in the following aspects:

1. Resolution is very poor due to the low solubility of Fe-Metal complexes in DMSO.
2. The disappearance of the signal due to the imine group, is attributed to its involvement in coordinating the Iron ion ^[19].

3. The spectrum displays a multiple signal at δ (6.58–7.63 ppm) assigned to aromatic ring protons.

The ^1H NMR Spectra of metal complexes are shown in Figure 7, 8, 9.




Fig 7: ^1H NMR of tris(2-((E)-1-hydrazoneethyl)-4,5-dimethylphenoxy)iron




Fig 8: ^1H NMR of tris(4,5-dimethyl-2-((E)-1-(2-phenylhydrazone)ethyl) phenoxy)iron.




Fig 9: ^1H NMR of tris (2-((E)-1-(2-(2,4-dinitrophenyl)hydrazono)ethyl)-4,5-dimethyl phenoxy) iron

Conclusion

All Fe complexes are characterized by elemental analysis, IR, UV Visible spectrophotometer, ^1H NMR, Molar Conductance and Magnetic susceptibility measurement, gravimetric analysis. Based on above results it suggests the Octahedral geometry for the Fe (III) complexes and the conductivity data show that each one these complexes are non-electrolytes.

Acknowledgements

The authors thank Mr. Pravin Daule and Mr. Dhole Anil, PG Department of Chemistry, New Arts, Commerce and Science College Parner, University of Pune, Dr. Raj Badekar, Riva Industries Ambernath, Mumbai, Dr. Pradeep Swain, Aveum Biolabs Pvt. Ltd., Hyderabad and Department of chemistry and Institute of Science, Mumbai for their valuable guidance and providing necessary facilities.

References

1. Trehoux J, Mahy P, Avenier F. A growing family of O₂ activating dinuclear iron enzymes with key catalytic diiron(III)-peroxo intermediates: Biological systems and chemical models. *Coordination Chemistry Reviews*. 2016; 322:142-158.
2. Jasniewski AJ, Que Jr L. Dioxygen Activation by Nonheme Diiron Enzymes: Diverse Dioxygen Adducts, High-Valent

Intermediates, and Related Model Complexes. *Chem. Rev.* 2018; 118:2554-2592.

3. Feig AL, Lippard SJ. Reactions of Non-Heme Iron (II) Centers with Dioxygen in Biology and Chemistry. *Chem. Rev.* 1994; 94:759-805.
4. Torre MH, Gambino D, Araujo J, Cerecetto H, Gonza^lez M, Azqueta ML, *et al.* Novel Cu(II) quinoxaline N1, N4-dioxide complexes as selective hypoxic cytotoxins. *European Journal of Medicinal Chemistry*. 2005; 40:473-480.
5. Recharadson DR, Bernhardt PV. Crystal and molecular structure of 2-hydroxy-1-naphthaldehyde isonicotinoyl hydrazone (NIH) and its iron (III) complex: an iron chelator with anti-tumour activity. *Journal of Biological Inorganic Chemistry*. 1999; 4:266-273.
6. Brookhart M, Small BL. Polymerization of Propylene by a New Generation of Iron Catalysts: Mechanisms of Chain Initiation, Propagation, and Termination. *Macromolecules*. 1999; 32:2120-2130.
7. Flavia N, Angela L, Giancarlo M, Carlo P, Vincenzo P, Genevieve C, *et al.* Hemoprotein models based on a covalent helix-heme-helix sandwich. 3. Coordination properties, reactivity and catalytic application of Fe (III)- and Fe(II)-mimo chrome I. *Journal of Biological Inorganic Chemistry*. 1998; 3:671-681.
8. Patel KN, Patel KM, Patel MN, Patel CR, Patel NH. Synthesis, Characterization and Antimicrobial Activities of Some Mixed-Ligand Complexes. *Synth React Inorg Met-Org Chem*. 2000; 30:1617-1627.

9. Routier S, Vezin H, Lamour E, Bernier JL, Cattean JP, Bailly C, *et al.* DNA cleavage by hydroxy-salicylidene-ethylendiamine-iron complexes. *Nucleic Acid Research*. 1999; 27:4160-4166.
10. Catherinr H, Marguerite P, Michael R, Heinz G, Ste'phanie S, Bernard M, *et al.* Preparation, characterization and crystal structures of manganese(II), iron(III) and copper(II) complexes of the bis[di-1,1-(2-pyridyl)ethyl]amine (BDPEA) ligand; evaluation of their DNA cleavage activities. *Journal of Biological Inorganic Chemistry*. 2001; 6:14-22.
11. Yamgar RS, Nivid YR, Nalawade S, Mandewale M, Atram RG, Sawant Novel Zinc (II) SS. *Complexes of Heterocyclic Ligands as Antimicrobial Agents: Synthesis, Characterisation, and Antimicrobial Studies*. *Bioinorganic Chemistry and Applications*. 2014; (Article id 76598): 10.
12. Jayabalakrishnan C, Natarajan K. Synthesis, characterization, and biological activities of ruthenium (II) carbonyl complexes containing bifunctional tridentate Schiff bases. *Synthesis and Reactivity in Inorganic and Metal-Organic Chemistry*. 2001; 31:983-995.
13. Mandewale MC, Thorat BR, Shelke D, Patil R, Yamgar R. Synthesis, characterization and fluorescence study of N-(E)-(2-hydroxyquinolin-3-yl) methylidene]-1-benzofuran- 2-carbohydrazide and its metal complexes. *Heterocyclic letters*. 2015; 5:251-259.
14. Dharmaraj N, Viswanathamurthi P, Natarajan K. Ruthenium (II) complexes containing bidentate Schiff bases and their antifungal activity. *Transition Metal Chemistry*. 2001; 26:105-109.
15. Mandewale MC, Thorat BR, Yamgar RS. Synthesis and anti-mycobacterium study of some fluorine containing Schiff bases of quinoline and their metal complexes. *Der Pharma Chemica*. 2015; 7:207-215.
16. Dinkar Gagare, Raju Patil, Rama Lokhande, Poonam Dwivedi, Ajay Deshmukh. Synthesis and characterization of (E)-2-(1-hydrazonoethyl)-4, 5-dimethylphenol from 2-hydroxy-4, 5-dimethylacetophenone. *J. Chem. Sci.* 2020; 132:2-8.
17. Dinkar Gagare, Raju Patil, Rama Lokhande, Poonam Dwivedi, Ajay Deshmukh. Synthesis and characterization of (E)-2-(1-(2-(2,4-dinitrophenyl)hydrazono)ethyl)-4,5-dimethylphenol from 2-hydroxy-4,5-dimethylacetophenone. *International Journal of Advanced Chemistry Research*. 2020; 2:36-40.
18. Dinkar Gagare, Raju Patil, Rama Lokhande, Poonam Dwivedi, Ajay Deshmukh. Synthesis and characterization of (E)-4, 5-dimethyl-2-(1-(2-phenylhydrazono) ethyl) phenol from 2-hydroxy-4, 5-dimethylacetophenone. *International Journal of Advanced Chemistry Research*. 2020; 2: 41-47.
19. Nora H, Al-Sha'alan. Antimicrobial activity and spectral, magnetic and thermal studies of some transition metal complexes of a Schiff base hydrazone containing a quinoline moiety. *Molecules*. 2007; 12:1080-1091.