



## Effect of FYM, Zn and Fe fertilizer on Physio-chemical properties of soil under Pearlmillet in hyper arid region of Rajasthan

Basu Devi Yadav<sup>1\*</sup>, SR Yadav<sup>2</sup>, PS Shekhawat<sup>1</sup>, Sushil Kumar Kharia<sup>1</sup>, Rajendra Kumar Yadav<sup>3</sup> and SL Yadav<sup>3</sup>

<sup>1</sup>College of Agriculture, Swami Keshwanand Rajasthan Agricultural University, Bikaner, Rajasthan, India

<sup>2</sup>Agricultural Research Station, Swami Keshwanand Rajasthan Agricultural University, Bikaner, Rajasthan, India

<sup>3</sup>Agricultural Research Station, Agricultural University, Kota, Rajasthan, India

DOI: <https://doi.org/10.33545/26646781.2021.v3.i2a.67>

### Abstract

A two years field experiment was conducted at two experimental sites of loamy sand soil. During *Kharif*, 2019 experiment was conducted at Agricultural Research Farm and during *Kharif*, 2020 conducted at Instructional farm, COA, SKRAU, Bikaner. To evaluate the influence of application of FYM, Zn and Fe fertilizer on physio-chemical properties of soil after harvest of pearl millet (RHB-177). Treatments consisted of three levels of FYM (Control, FYM @ 5 t ha<sup>-1</sup> and 10 t ha<sup>-1</sup> FYM) and ten methods and levels of Zn and Fe application (Control, soil application of (25 kg ha<sup>-1</sup> FeSO<sub>4</sub>, 25 kg ha<sup>-1</sup> ZnSO<sub>4</sub>, 25 kg ha<sup>-1</sup> ZnSO<sub>4</sub> + ZSB, 25 kg ha<sup>-1</sup> FeSO<sub>4</sub> + 25 kg ha<sup>-1</sup> ZnSO<sub>4</sub> + ZSB) and foliar application of (0.5% FeSO<sub>4</sub>, 1.0% FeSO<sub>4</sub>, 0.5% ZnSO<sub>4</sub>, 0.5% FeSO<sub>4</sub> + 0.5% ZnSO<sub>4</sub>, 0.5% FeSO<sub>4</sub> + 0.5% ZnSO<sub>4</sub>) in split plot design with three replications. The results revealed that organic carbon, dehydrogenase activity, bacterial, fungal and actinomycetes count, available nitrogen, phosphorus, potassium and sulphur in soil after harvest of pearl millet was significantly increased, while saturated hydraulic conductivity was significantly decreased with increasing levels of FYM up to @ 10 t ha<sup>-1</sup> over control and 5 t FYM ha<sup>-1</sup>. And application of different methods and levels of Zn and Fe manifested significantly positive impact on available sulphur, whereas available phosphorus was significantly decreased as compare to control.

**Keywords:** FYM, Zn and Fe, pearlmillet, physio-chemical properties of soil

### Introduction

Pearl millet (*Pennisetum glaucum* (L.) R. Br. emend Stuntz) is the most widely grown drought tolerant warm season coarse grain cereal and is considered as a Poor man's food crop. Pearl millet is one of the fourth most important crop in India after rice, wheat and sorghum which flourishes well even under adverse conditions of weather, low soil fertility and alkaline soils (Singh *et al.*, 2017; Shekhawat and Kumawat, 2017) <sup>[7, 16]</sup>. Pearl millet is provided nutritional security as it has higher protein content (12.6%), more balanced amino acid profile, and contribute about one third of iron and zinc requirements (Manga and Kumar, 2011) <sup>[10]</sup>. In Rajasthan, Pearl millet is mostly cultivated as unirrigated crop, unfertilized or partially fertilized on marginal and submarginal lands having light textured soils, low in nitrogen, phosphorus and organic matter with poor moisture retention capacity, thereby resulting in low and uneconomical harvest. Hence, our research effort should be diverted to remove the constraints that are responsible for its poor yield. Intense temperature and low organic matter content in semi-arid region leads to poor soil fertility and low nitrogen status (Rego *et al.*, 2003) <sup>[15]</sup>. Application of FYM to crops also avoids its wasteful use for burning purposes. FYM is being used as a major source of organic manure in field crops. Limited availability of this manure is, however, an important constraint in its use as a source of nutrients. FYM is rich in organic matter and is a good source of plant nutrients. Poor nutrient economy of light textured soil necessitates the need for supplementing fertilizers with FYM. Organic manure influence both yield and plant micronutrients need and thus help to sustain crop productivity. Approximately 49 per cent of world & 50 per cent of Indian soils are deficient in Zn (Cakmak *et al.*, 2010; Shekhawat and Kumawat, 2017) <sup>[2, 16]</sup>. Continuous intensive

cropping of high yielding varieties has further aggravated the depletion of soil zinc leading to low zinc concentration in edible grains. The soils of hyper arid region are coarse textured and deficient in N, S, Zn and Fe due to continuous crops removal.

FYM changes the chemical, physical and biological properties of the soil which in terms affect the availability of plant nutrients. Soil application of ZnSO<sub>4</sub> and FeSO<sub>4</sub> each @ 25 kg ha<sup>-1</sup> increase the availability of S, Zn and Fe. Hence, keeping all these facts, the present study was undertaken to assessment of physio-chemical properties of soil through application of FYM, Zn and Fe fertilizer under pearlmillet in hyper arid region of Rajasthan.

### Material and Methods

During *kharif*, 2019, the experiment was conducted at experimental farm, Agriculture Research Station, SKRAU, Bikaner, and in *kharif*, 2020, the experiment was conducted at instruction farm, College of Agriculture, SKRAU, Bikaner. These sites were situated at 234 meters above mean sea level on latitude 28°10' N and 73°22' E longitude. Bikaner has arid climate with average annual rainfall of about 262 mm. More than 80 per cent of rainfall is received during *kharif* season (July-September) by the South West monsoon. During summer, the maximum temperature may go as high as 48° C while in the winters it may fall as low as 0°C. The soil is loamy sand desert soils with low water holding capacity, slightly alkaline in reaction (pH 8.24), poor in organic carbon (0.13 per cent), low in available nitrogen (104.2 kg ha<sup>-1</sup>) and medium in available phosphorus (16.68 kg ha<sup>-1</sup>) and medium in available potassium (178.05 kg ha<sup>-1</sup>). Similarly, *kharif*, 2020 location analytical results revealed that the soil of the experimental field was sandy in texture and slightly alkaline in

reaction (pH 8.29), poor in organic carbon (0.12 per cent), low in available nitrogen (101.36 kg ha<sup>-1</sup>) and medium in available phosphorus (17.12 kg ha<sup>-1</sup>) and medium in available potassium (182.32 kg ha<sup>-1</sup>).

Treatments consisted of three levels of FYM (Control, FYM @ 5 t ha<sup>-1</sup> and 10 t ha<sup>-1</sup> FYM) and ten methods and levels of Zn and Fe application (Control, soil application of (25 kg ha<sup>-1</sup> FeSO<sub>4</sub>, 25 kg ha<sup>-1</sup> ZnSO<sub>4</sub>, 25 kg ha<sup>-1</sup> ZnSO<sub>4</sub> + ZSB, 25 kg ha<sup>-1</sup> FeSO<sub>4</sub> + 25 kg ha<sup>-1</sup> ZnSO<sub>4</sub> + ZSB) and foliar application of (0.5% FeSO<sub>4</sub>, 1% FeSO<sub>4</sub>, 0.5% ZnSO<sub>4</sub>, 0.5% FeSO<sub>4</sub> + 0.5% ZnSO<sub>4</sub>, 0.5% FeSO<sub>4</sub> + 0.5% ZnSO<sub>4</sub>) in split plot design with three replications were applied to pearl millet crop (RHB-177). FYM was applied at 21 days before sowing @ 5 t ha<sup>-1</sup> and 10 t ha<sup>-1</sup> in the field as per treatment. RDF was given through Urea (N), SSP (P) and MOP (K) in each plot. Soil application of Zn and Fe were applied as per treatments through zinc sulphate (ZnSO<sub>4</sub>.7H<sub>2</sub>O) containing 21% Zn and 10% S and iron sulphate (FeSO<sub>4</sub>.7H<sub>2</sub>O) containing 19.6% Fe and 10.5% S at the time of sowing as basal dose and foliar application of zinc (0.5%) and iron sulphate (0.5% and 1%) at 25 and 40 DAS.

## Results and Discussions

### Soil physical properties

#### Effect of FYM, Zn and Fe on bulk density, particle density and saturated hydraulic conductivity

Effect of FYM, Zn and Fe application were found non-significant on bulk density and particle density of soil. Saturated hydraulic conductivity of soil significantly decreased due to FYM application. Saturated hydraulic conductivity of soil was varied from 12.16 to 10.96 cm hr<sup>-1</sup> on the pooled mean basis (Table 1). The lowest hydraulic conductivity was recorded with application of FYM @ 10 t ha<sup>-1</sup> (10.91, 11.1 and 10.96 cm hr<sup>-1</sup>), while highest saturated hydraulic conductivity was recorded in controlled plot (12.04, 12.28 and 12.16) during both years and pooled data, respectively. It could be due to granulation effect of farmyard manure which led to increase in soil micro porosity and thus reduction in water movement through soil. These findings are in agreement with earlier findings by Kumawat, 2019 [9]; Kumar, 2021 [8]. And non-significantly influenced due to methods and levels of Zn and Fe application.

### Biological properties

#### Effect of methods & levels of Zn, Fe and FYM on dehydrogenase activity

Different levels of FYM significantly increased dehydrogenase activity (Table 2) in soil after harvest of pearl millet over control. Significantly highest (24.38 µg TPF g<sup>-1</sup> soil 24 h<sup>-1</sup>) dehydrogenase activity was recorded with application of FYM @ 10 t ha<sup>-1</sup> which was significantly higher than FYM @ 5 t ha<sup>-1</sup> (21.96 µg TPF g<sup>-1</sup> soil 24 h<sup>-1</sup>) and control (21.15 µg TPF g<sup>-1</sup> soil 24 h<sup>-1</sup>) during both years and pooled data. On the basis of pooled data FYM @ 10 t ha<sup>-1</sup> increased dehydrogenase activity in the tune of 15.27 per cent over control. The dehydrogenase activity and microbial count did not influence significantly due to methods and levels of Zn and Fe application.

#### Effect of methods & levels of Zn, Fe and FYM on bacterial counts, fungi counts and actinomycetes counts

The results pertaining to bacterial count in soil presented in (Table 2) after harvest of pearl millet in both the years and as pooled mean basis indicated that bacterial count was varied from 20.85 to 28.10 × 10<sup>6</sup>. Application of different levels of FYM increase bacterial count in soil. Significantly highest bacterial count was recorded with application of FYM @ 10 t

ha<sup>-1</sup>, while lowest bacterial count was recorded in controlled condition and superior over 5 t ha<sup>-1</sup> FYM.

And fungi count was varied from 15.62 to 24.60 × 10<sup>4</sup>. Application of different levels of FYM increase fungi count in soil. Significantly highest fungi count was recorded with application of FYM @ 10 t ha<sup>-1</sup> (24.60 × 10<sup>4</sup>), while lowest bacterial fungi count was recorded in controlled condition (15.62 × 10<sup>4</sup>) and superior over 5 t ha<sup>-1</sup> FYM (19.17 × 10<sup>4</sup>) in pooled mean basis.

FYM significantly increased actinomycetes count (Table 2) in soil after harvest of pearl millet over control during both years. Significantly highest (43.69 × 10<sup>5</sup>) actinomycetes count was recorded with application of FYM @ 10 t ha<sup>-1</sup> which was significantly higher than FYM @ 5 t ha<sup>-1</sup> (40.74 × 10<sup>5</sup>) and control (37.90 × 10<sup>5</sup>) on the basis of pooled mean. The microbial count did not influence significantly due to methods and levels of Zn and Fe application. The data showed that application of FYM levels significantly increased all biological properties as compared to control. It due to FYM act as source of organic carbon and energy, which stimulate different microbial activities in soil (Bhatt *et al.*, 2016; Rajput *et al.*, 2019; Kumar, 2021) [1, 14, 8]. The increase in microbial count might be due to increasing levels of N, P, K, S, Zn and Fe which might have increased cation exchange capacity, root biomass, root exudates and ultimately provided carbon and energy to the soil microbes resulting into multiplication of microbial count. The similar results were also reported by Parewa *et al.*, 2014 [12], Urmila *et al.*, 2016 [19]; Dudhawal, 2018.

### Chemical properties

#### Effect of FYM, Zn and Fe application on pH and electrical conductivity, organic carbon, available N, P, K and S

pH and electrical conductivity of soil did not influence significantly due to FYM levels and different methods and levels of Zn and Fe application.

Organic carbon of soil increased significantly (Table 3) due to application of different FYM levels during both the years as well as pooled analysis. Significantly the highest organic carbon (1.40 g kg<sup>-1</sup>) content was observed with application of FYM @ 10 t ha<sup>-1</sup> as compared to control (1.25 g kg<sup>-1</sup>) and FYM @ 5 t ha<sup>-1</sup> (1.29 g kg<sup>-1</sup>) on pooled mean basis.

Available nitrogen (Table 4) in soil after harvest of pearl millet during both the years and pooled analysis increased significantly due to application of different FYM levels. It was varied from 101.9 to 110.9 kg ha<sup>-1</sup>. Significantly the highest available nitrogen (110.9 kg ha<sup>-1</sup>) was recorded with application of FYM @ 10 t ha<sup>-1</sup> which was statistically higher than FYM @ 5 t ha<sup>-1</sup> and control.

Available phosphorous in soil after harvest of pearl millet during both the years and pooled analysis increased significantly (Table 4) due to application of different FYM levels. It was varied from 15.22 to 17.51 kg ha<sup>-1</sup>. Significantly the highest available phosphorous (17.50 kg ha<sup>-1</sup>) was recorded with application of FYM @ 10 t ha<sup>-1</sup> which was statistically higher than FYM @ 5 t ha<sup>-1</sup> and control. And application of different methods and levels of Zn and Fe significantly decreased available phosphorous. All treatments of soil application of ZnSO<sub>4</sub> had significantly lower (Table 4) available phosphorous than control. On the basis of pooled data soil application of ZnSO<sub>4</sub> @ 25 kg ha<sup>-1</sup> was significantly decrease available phosphorous (15.15 kg ha<sup>-1</sup>) than control (16.59 kg ha<sup>-1</sup>).

Available potassium in soil after harvest of pearl millet during

both the years and pooled analysis increased significantly (Table 4) due to application of different FYM levels. It was varied from 175.0 to 188.7 kg ha<sup>-1</sup>. Significantly the highest available potassium (188.7 kg ha<sup>-1</sup>) was recorded with application of FYM @ 10 t ha<sup>-1</sup> which was statistically higher than FYM @ 5 t ha<sup>-1</sup> (181.9 kg ha<sup>-1</sup>) and control (175.0 kg ha<sup>-1</sup>) on pooled data basis.

Progressive increase in levels of FYM up to 10 t ha<sup>-1</sup> significantly increased available sulphur in soil after harvest of pearl millet over control (Table 4) during both the individual years and pooled analysis. Application of FYM @ 10 t ha<sup>-1</sup> significantly influenced (16.28 kg ha<sup>-1</sup>) available sulphur compared to FYM @ 5 t ha<sup>-1</sup> (14.98 kg ha<sup>-1</sup>) and control (13.58 kg ha<sup>-1</sup>). Sulphur was significantly increased with application of different methods and levels of Zn and Fe over control. All treatments of soil application of ZnSO<sub>4</sub> and FeSO<sub>4</sub> had significantly higher available sulphur than control during both years of study. On the basis of pooled data soil application of FeSO<sub>4</sub> @ 25 kg ha<sup>-1</sup>+ ZnSO<sub>4</sub> @ 25 kg ha<sup>-1</sup>+ZSB was significantly increase available sulphur (19.27 kg ha<sup>-1</sup>) than control (13.44 kg ha<sup>-1</sup>). And This might be due to adsorption of part of applied sulphur on organic matter and thereby reducing the leaching losses of sulphur.

The use of FYM through proper practice helps in increasing the organic carbon content of soil due to higher biomass and differential rate of oxidation of organic matter by microbes. The similar effects were also observed by (Parewa *et al.*, 2014 [12] and Choudhary *et al.*, 2019 [3]. Increases in availability of nitrogen, phosphorus, potassium and sulphur in soil under FYM treated plot might be due to FYM attributed to its inherent capacity to add good amount of organic carbon to the soil which hastens the process of mineralization of organically bound nitrogen and phosphorus present in native soil due to masking of adsorption sites by humate ions (Pattanayak *et al.*, 2009 and Kaur, 2013; Dudhawal, 2018) [13, 6]. And available phosphorus significantly decreased due to methods and levels of Zn and Fe application. It might be due to the antagonistic effect between Zn on P in soils solution and forming insoluble compounds, Zn<sub>3</sub>(PO<sub>4</sub>)<sub>2</sub> resulting in the low amount of P in the available pool. but non-significant difference was found with foliar application of Zn and Fe. Such types of findings were also reported by Keram *et al.*, 2012 [7] and Meena *et al.*, 2017 [11]. Higher available S in soil is due to greater mineralization of organic S and release of SO<sub>4</sub><sup>2-</sup> ions on its gradual oxidation. Similar findings were also reported by Tiwari *et al.* (1992) [18]; Jat *et al.* (2012) [5].

**Table 1:** Effect of FYM levels and methods & levels of Zn and Fe on bulk density (BD), particle density (PD) and saturated hydrolic conductivity (SHC) in soil after harvest of pearl millet

	Treatments	BD (Mg m <sup>-3</sup> )			PD (Mg m <sup>-3</sup> )			SHC (cm hr <sup>-1</sup> )		
		2019	2020	Pooled	2019	2020	Pooled	2019	2020	Pooled
<b>Farm yard manure levels</b>										
	M <sub>0</sub> -Control	1.53	1.53	1.53	2.61	2.62	2.62	12.04	12.28	12.16
	M <sub>1</sub> -5 t ha <sup>-1</sup>	1.51	1.51	1.51	2.60	2.61	2.60	11.48	11.69	11.58
	M <sub>2</sub> -10 t ha <sup>-1</sup>	1.50	1.48	1.49	2.59	2.59	2.59	10.91	11.01	10.96
	S.Em±	0.01	0.02	0.01	0.02	0.03	0.02	0.10	0.07	0.06
	CD at 5%	NS	NS	NS	NS	NS	NS	0.40	0.26	0.20
<b>Methods and Levels of Zn &amp; Fe application</b>										
Soil application	F <sub>0</sub> -Control	1.52	1.53	1.52	2.61	2.62	2.61	11.63	11.91	11.77
	F <sub>1</sub> -25 kg ha <sup>-1</sup> FS	1.52	1.51	1.51	2.60	2.61	2.60	11.59	11.71	11.65
	F <sub>2</sub> -25 kg ha <sup>-1</sup> ZS	1.50	1.51	1.50	2.61	2.59	2.60	11.61	11.76	11.68
	F <sub>3</sub> - 25 kg ha <sup>-1</sup> ZS + ZSB	1.51	1.49	1.50	2.60	2.61	2.60	11.39	11.61	11.50
	F <sub>4</sub> -25 kg ha <sup>-1</sup> FS + 25 kg ha <sup>-1</sup> ZS + ZSB	1.49	1.50	1.49	2.59	2.59	2.59	11.19	11.65	11.42
Foliar application	F <sub>5</sub> -0.5% FS	1.52	1.49	1.51	2.60	2.62	2.61	11.49	11.60	11.54
	F <sub>6</sub> -1.0% FS	1.51	1.51	1.51	2.61	2.60	2.61	11.44	11.61	11.52
	F <sub>7</sub> -0.5% ZS	1.52	1.50	1.51	2.60	2.61	2.60	11.41	11.54	11.48
	F <sub>8</sub> -0.5% FS+0.5% ZS	1.52	1.52	1.52	2.61	2.60	2.61	11.51	11.61	11.56
	F <sub>9</sub> -1.0% FS+0.5% ZS	1.51	1.51	1.51	2.59	2.61	2.60	11.49	11.59	11.54
	S.Em±	0.02	0.02	0.01	0.04	0.03	0.03	0.17	0.08	0.09
	CD at 5%	NS	NS	NS	NS	NS	NS	NS	NS	NS

FS= Ferrous Sulphate, ZS= Zinc Sulphate, ZSB= Zinc Solubilizing Bacteria

**Table 2:** Effect of FYM levels and methods & levels of Zn and Fe on biological properties of post-harvest soil under pearl millet

	Treatments	Dehydrogenase (Ug TPF g <sup>-1</sup> soil 24 h <sup>-1</sup> )			Bacteria Count (10 <sup>6</sup> )			Fungi Count (10 <sup>4</sup> )			Actinomycetes Count (10 <sup>5</sup> )		
		2019	2020	Pooled	2019	2020	Pooled	2019	2020	Pooled	2019	2020	Pooled
<b>Farm yard manure levels</b>													
	M <sub>0</sub> -Control	21.71	20.59	21.15	21.26	20.45	20.85	16.36	14.89	15.62	38.33	37.47	37.90
	M <sub>1</sub> -5 t ha <sup>-1</sup>	22.01	21.90	21.96	24.83	21.92	23.37	22.05	17.38	19.71	41.68	39.79	40.74
	M <sub>2</sub> -10 t ha <sup>-1</sup>	24.22	24.55	24.38	28.64	27.56	28.10	25.35	23.84	24.60	43.78	43.60	43.69
	S.Em±	0.19	0.17	0.13	0.19	0.23	0.15	0.15	0.15	0.10	0.37	0.17	0.20
	CD at 5%	0.74	0.68	0.42	0.74	0.89	0.48	0.57	0.59	0.34	1.46	0.66	0.66
<b>Methods and Levels of Zn &amp; Fe application</b>													
Soil application	F <sub>0</sub> -Control	22.1	22.2	22.2	24.4	23.0	23.7	21.2	18.5	19.8	40.2	40.0	40.1
	F <sub>1</sub> -25 kg ha <sup>-1</sup> FS	22.75	22.63	22.69	24.95	23.44	24.20	21.24	18.60	19.92	41.43	40.50	40.97
	F <sub>2</sub> -25 kg ha <sup>-1</sup> ZS	22.77	22.65	22.71	24.84	23.45	24.15	21.24	18.71	19.97	41.48	40.60	41.04
	F <sub>3</sub> - 25 kg ha <sup>-1</sup> ZS + ZSB	22.80	22.65	22.72	25.03	23.52	24.27	21.28	18.77	20.03	41.56	40.62	41.09
	F <sub>4</sub> -25 kg ha <sup>-1</sup> FS + 25 kg ha <sup>-1</sup> ZS + ZSB	22.82	22.71	22.76	25.09	23.49	24.29	21.52	19.13	20.32	41.60	40.69	41.15
Foliar application	F <sub>5</sub> -0.5% FS	22.42	22.07	22.24	24.93	23.23	24.08	21.19	18.51	19.85	41.01	40.02	40.52
	F <sub>6</sub> -1.0% FS	22.52	22.21	22.37	25.07	23.38	24.22	21.20	18.72	19.96	41.15	40.10	40.62

	F7-0.5% ZS	22.61	22.34	22.48	25.01	23.20	24.11	21.21	18.70	19.96	41.43	40.12	40.78
	F8-0.5% FS+0.5% ZS	22.86	21.94	22.40	24.85	23.36	24.11	21.22	18.85	20.04	41.36	40.04	40.70
	F9-1.0% FS+0.5% ZS	22.77	22.02	22.40	24.87	23.02	23.95	21.23	18.52	19.88	41.44	40.15	40.80
	S.Em±	0.29	0.30	0.21	0.25	0.34	0.21	0.22	0.29	0.18	0.52	0.59	0.39
	CD at 5%	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS

FS= Ferrous Sulphate, ZS= Zinc Sulphate, ZSB= Zinc Solubilizing Bacteria

**Table 3:** Effect of FYM levels and methods & levels of Zn and Fe on pH, electric conductivity (EC), organic carbon (OC) in soil in soil after harvest of pearl millet

	Treatments	pH			EC (dSm <sup>-1</sup> )			OC (g kg <sup>-1</sup> )		
		2019	2020	Pooled	2019	2020	Pooled	2019	2020	Pooled
<b>Farm yard manure levels</b>										
	M0-Control	8.22	8.26	8.24	0.171	0.182	0.176	1.31	1.20	1.25
	M1-5 t ha <sup>-1</sup>	8.11	8.20	8.15	0.172	0.183	0.178	1.35	1.22	1.29
	M2-10 t ha <sup>-1</sup>	8.09	8.14	8.11	0.174	0.186	0.180	1.45	1.36	1.40
	S.Em±	0.06	0.05	0.04	0.002	0.002	0.001	0.01	0.01	0.01
	CD at 5%	NS	NS	NS	NS	NS	NS	0.06	0.04	0.03
<b>Methods and Levels of Zn &amp; Fe application</b>										
Soil application	F0-Control	8.20	8.26	8.23	0.171	0.182	0.177	1.36	1.24	1.30
	F1-25 kg ha <sup>-1</sup> FS	8.21	8.25	8.23	0.178	0.184	0.181	1.37	1.28	1.33
	F2-25 kg ha <sup>-1</sup> ZS	8.16	8.17	8.17	0.172	0.184	0.178	1.38	1.25	1.32
	F3- 25 kg ha <sup>-1</sup> ZS + ZSB	8.07	8.08	8.08	0.176	0.186	0.181	1.39	1.28	1.34
	F4-25 kg ha <sup>-1</sup> FS + 25 kg ha <sup>-1</sup> ZS + ZSB	7.98	8.14	8.06	0.176	0.186	0.181	1.39	1.29	1.34
Soil application	F5-0.5% FS	8.18	8.24	8.21	0.168	0.183	0.175	1.36	1.24	1.30
	F6-1.0% FS	8.15	8.24	8.19	0.170	0.183	0.176	1.36	1.26	1.31
	F7-0.5% ZS	8.18	8.25	8.22	0.174	0.185	0.179	1.37	1.24	1.31
	F8-0.5% FS+0.5% ZS	8.15	8.18	8.17	0.172	0.183	0.178	1.36	1.26	1.31
	F9-1.0% FS+0.5% ZS	8.12	8.17	8.15	0.168	0.179	0.174	1.37	1.25	1.31
	S.Em±	0.09	0.08	0.06	0.003	0.002	0.002	0.02	0.02	0.01
	CD at 5%	NS	NS	NS	NS	NS	NS	NS	NS	NS

FS= Ferrous Sulphate, ZS= Zinc Sulphate, ZSB= Zinc Solubilizing Bacteria

**Table 4:** Effect of FYM levels and methods & levels of Zn and Fe on available nitrogen, available phosphorous, available potassium and Available sulphur in soil after harvest of pearl millet

	Treatments	Avail. Nitrogen (Kg ha <sup>-1</sup> )			Avail. phosphorous (Kg ha <sup>-1</sup> )			Avail. potassium (Kg ha <sup>-1</sup> )			Avail. sulphur (Kg ha <sup>-1</sup> )		
		2019	2020	Pooled	2019	2020	Pooled	2019	2020	Pooled	2019	2020	Pooled
<b>Farm yard manure levels</b>													
	M0-Control	102.9	100.9	101.9	15.49	14.95	15.22	173.7	176.3	175.0	14.67	12.49	13.58
	M1-5 t ha <sup>-1</sup>	109.7	105.7	107.7	16.75	16.00	16.37	180.7	183.2	181.9	15.59	14.38	14.98
	M2-10 t ha <sup>-1</sup>	113.8	107.9	110.9	18.11	16.91	17.51	187.6	189.9	188.7	17.29	15.26	16.28
	S.Em±	0.59	0.68	0.45	0.16	0.16	0.11	1.60	1.61	1.13	0.15	0.14	0.10
	CD at 5%	2.31	2.66	1.46	0.62	0.62	0.36	6.27	6.31	3.69	0.57	0.56	0.33
<b>Methods and Levels of Zn &amp; Fe application</b>													
Soil application	F0-Control	107.0	103.3	105.2	16.96	16.21	16.59	178.4	181.2	179.8	14.74	12.15	13.44
	F1-25 kg ha <sup>-1</sup> FS	109.1	104.9	107.0	17.56	16.59	17.08	179.6	183.6	181.6	15.90	15.10	15.50
	F2-25 kg ha <sup>-1</sup> ZS	109.5	105.6	107.5	15.68	14.63	15.15	181.3	185.1	183.2	16.41	15.31	15.86
	F3- 25 kg ha <sup>-1</sup> ZS + ZSB	109.9	105.1	107.5	16.07	14.72	15.40	183.2	185.6	184.4	16.90	16.41	16.65
	F4-25 kg ha <sup>-1</sup> FS + 25 kg ha <sup>-1</sup> ZS + ZSB	110.0	105.0	107.5	16.10	15.06	15.58	183.3	185.8	184.6	19.97	18.57	19.27
Foliar application	F5-0.5% FS	108.8	104.3	106.6	17.26	16.67	16.97	178.4	181.1	179.7	14.88	12.55	13.72
	F6-1.0% FS	107.4	104.6	106.0	16.98	16.65	16.82	179.3	181.6	180.4	14.92	12.72	13.82
	F7-0.5% ZS	108.6	104.2	106.4	17.29	16.41	16.85	179.8	183.9	181.8	14.89	12.60	13.75
	F8-0.5% FS+0.5% ZS	108.9	105.6	107.2	16.98	16.26	16.62	180.8	182.1	181.4	15.01	12.66	13.83
	F9-1.0% FS+0.5% ZS	108.6	105.8	107.2	16.91	16.33	16.62	182.4	181.4	181.9	14.88	12.38	13.63
	S.Em±	1.23	1.22	0.86	0.25	0.26	0.18	2.85	2.85	2.02	0.25	0.21	0.16
	CD at 5%	NS	NS	NS	0.71	0.72	0.50	NS	NS	NS	0.70	0.58	0.45

FS= Ferrous Sulphate, ZS= Zinc Sulphate, ZSB= Zinc Solubilizing Bacteria

## Conclusion

Result of the present study shows that significant improvement in organic carbon, available N, P, K and S, biological parameters (Bacterial count, fungal count, actinomycetes and dehydrogenase activity) whereas, saturated hydraulic conductivity significantly decreased with application of FYM @ 10 t ha<sup>-1</sup>. Available sulphur significantly increased with Zn and Fe application.

## References

- Bhatt B, Chandra R, Ram S, Pareek N. Long-term effects of fertilization and manuring on productivity and soil biological properties under rice (*Oryza sativa*)-wheat (*Triticum aestivum*) sequence in mollisols. Archives of Agronomy and Soil Science. 2016;62:1109-1122.
- Cakmak I Pfeiffer, Wolfgang H, McClafferty B. Biofortification of durum wheat with zinc and iron. Cereal chemistry. 2010;87:10-20.

3. Choudhary A, Yadav SR, Parewa HP. Effect of wool waste in combination with farm yard manure and fertilizer on soil properties in arid soil of Bikaner. *Indian Journal of Environmental Biology*. 2019;40:1067-1072.
4. Dhudhawal A. Effect of soil management practices on soil properties and yield of pearl millet (*Pennisetum glaucum* (L.) R. Br. emend. Stuntz). M.Sc. Thesis, S K N Agricultural University, Jobner. 2018.
5. Jat G, Sharma KK, Jat NK. Effect of FYM and mineral nutrients on physico-chemical properties of soil under mustard in western arid zone of India. *Annals of Plant and Soil Research*. 2012;14:167-166.
6. Kaur S. Long-term effects of inorganic and integrated fertilization on soil phosphorus dynamics and crop uptake. M. Sc. Thesis, Punjab Agricultural University, Ludhiana, 2013.
7. Keram KS, Sharma BL, Sawarkar SD. Impact of Zn application on yield, quality, nutrients uptake and soil fertility in a medium deep black soil (vertisol). *International Journal of Science, Environment and Technology*. 2012;1:563-571.
8. Kumar S. Nutrient dynamics under wool waste based nutrient management in bottle gourd-onion cropping system in torripsamments of western rajasthan. Ph.D. thesis, S K Rajasthan Agricultural University, Bikaner. 2021.
9. Kumawat S. Impact of STCR approach on soil carbon fractions and productivity of clusterbean under clusterbean-wheat cropping system. M.Sc. thesis, S K Rajasthan Agricultural University, Bikaner, 2019.
10. Manga VK, Kumar A. Cultivar options for increasing pearl millet productivity in arid regions. *Indian Journal of Fundamental and Applied Life Sciences*. 2011;1:200-208.
11. Meena NR, Meena MK, Sharma KK, Meena MD. Effect of zinc enriched farm yard manure on yield of mungbean and physico-chemical properties of soil. *Legume Research*, 2017, LR-3929.
12. Parewa H, Yadav J, Rakshit A. Effect of Fertilizer Levels, FYM and Bioinoculants on Soil Properties in Inceptisol of Varanasi, Uttar Pradesh, India. *International Journal of Agriculture, Environment and Biotechnology*. 2014;7:517-525.
13. Pattanayak SK, Sureshkumar P, Tarafdar JC. New vista in phosphorus research. *Journal of the Indian Society of Soil Science*. 2009;57:536-545.
14. Rajput R, Pokhriya P, Panwar P, Arunchalam A, Arunchalam K. Soil nutrients, microbial biomass and crop response to organic amendments in rice cropping system in the Shiwaliks of Indian Himalayas. *International Journal of Recycling of Organic Waste in Agriculture*. 2019;8:73-85.
15. Rego TJ, Rao VN, Seeling B, Pardhasaradhi G, Kumar R. Nutrient balances-a guide to improving sorghum and groundnut-based dryland cropping systems in semi-arid tropical India. *Field Crops Research*. 2003;81:53-68.
16. Shekhawat PS, Kumawat N. Response of zinc fertilization on production and profitability of Pearl millet (*Pennisetum glaucum*) under rainfed condition of Rajasthan. *Journal of Agri. Search*. 2017;4:251-254.
17. Singh AK, Singh AK, Kumar R, Prakash V, Sundaram PK, Yadav SK. Indian Cereals Saga: Standpoint and Way Forward. *Journal of Agri Search*. 2017;4:1-10.
18. Tiwari RC, Singh SK, Pandey DK. Influence of sulphur application on yield and chemical composition of some crops. *Fertiliser News*. 1992;37:23-26.
19. Urmila Rawat US, Bhandari SC, Purohit HS. Studies on microbial indices of soil under long term nutrient management. *Ecology, Environment and Conservation*. 2016;22:2047-2051.