

ISSN Print: 2664-6781
 ISSN Online: 2664-679X
 Impact Factor: RJIF 5.32
 IJACR 2024; 6(1): 38-43
www.chemistryjournals.net
 Received: 11-12-2023
 Accepted: 19-01-2024

Zinah Talib Alsallami
 Department of Food Science,
 Faculty of Agriculture,
 University of Kufa, Najaf, Iraq

Ali Joodi Shuna Al-Fatlawi
 Department of Food Science,
 Faculty of Agriculture,
 University of Kufa, Najaf, Iraq

Hasan Najah Smesam
 Department of Chemistry,
 Faculty of Science, University
 of Kufa, Najaf, Iraq

**Murtadha Abdulhasan
 Aldhalemi**
 Department of Food Science,
 Faculty of Agriculture,
 University of Kufa, Najaf, Iraq

Qais R Lahhob
¹ Department of Food Science,
 Faculty of Agriculture,
 University of Kufa, Najaf, Iraq
² College of Pharmacy,
 National University of Science
 and Technology, Oman

Corresponding Author:
Qais R Lahhob
¹ Department of Food Science,
 Faculty of Agriculture,
 University of Kufa, Najaf, Iraq
² College of Pharmacy,
 National University of Science
 and Technology, Oman

The role of nutrition in maintaining the effectiveness of the liver (Enzymes and some blood variables) against heavy metals efficacy of leek powder in maintaining liver functions against cadmium negative effects

Zinah Talib Alsallami, Ali Joodi Shuna Al-Fatlawi, Hasan Najah Smesam, Murtadha Abdulhasan Aldhalemi and Qais R Lahhob

DOI: <https://doi.org/10.33545/26646781.2024.v6.i1.a.166>

Abstract

Interacting Cadmium as a critical pollution dilemma which includes human health risks currently in sight is one of the key issues in today's context because of the severe impacts to human health. Here, the experiment was aimed to specifically the effect of powder of leaves of leek plants on blood parameters and internal organs when toxicity effects of cadmium elements to rats were considered. The experimental groups consisted of a negative control (T₁: Option 1: For the experiment, T₁ was used as the control in which cadmium was injected to rats at a concentration of 0.4 µg/L, T₂ was chosen as the positive control with 25.7 µg/L of cadmium injected to rats, while T₃ and T₄ were the therapeutic interventions where rats were given 2% Results demonstrated that cadmium (T₂) significantly impacted blood cadmium levels (25.7 µg/L) compared to control (T₁: Concentrations of 2µm/L and higher were lethal (kidneys 5.4g/L, testes 5.1g/L, liver 19.1g/L, heart 0.8g/L and spleen 2.2g/L) and were damaging to internal organs compared to control (T₁: Kidneys 2.8 g/ As to Cadmium, we had increased enzyme levels in the blood at the conclusion compared to the control. AST was 22.5 U/l, ALT was 29.1 U/l and Alkaline Phosphatase was 18.3 IU/l in T₁, and in T₂, AST was 60.1 U/l, ALT was The pharmaceutical approach with leek was the best option compared to the others that gave the better outcomes which made the livers' functionality disrupted, blood cadmium levels, and internal organ weights. The article suggests leek, covering 2% of the dose, being a promising therapeutic agent in heavy metal cadmium detoxification. To date, such study is the first in the line of such researches that aims to demonstrate leek's presence as a component in cadmium toxicity remedy.

Keywords: Cadmium, leek, liver, kidneys, rat testicles

Introduction

Heavy metals are very often used in most various industrial, agricultural, and medical operations as well. Although the dominance of heavy metals over industries is considered priceless, these substances are an extremely critical phenomenon of danger and a great challenge for the companies. This project is focused on the habits of the certain personalities and the level of which they are performed in the present time and also the transport of metals in the human body through indirect actions of heavy metals with respect to their toxicity (Kang *et al.*, 2019; Yu *et al.*, 2020; Zou *et al.*, 2021) [6, 17, 20]. Cadmium is recognized as one of the mutually toxic heavy metals accumulated in human body (Hayat *et al.*, 2018; Wang *et al.*, 2017; Zheng *et al.*, 2020; Zou *et al.*, 2021) [3, 16, 19, 20]. On the whole, the individuals are exposed to the pollutants of cadmium anytime, as it is connected to inhaling gases, mining industries, plastic, ceramics and colored materials (Hayat *et al.*, 2018; Wang *et al.*, 2017; Zheng *et al.*, 2021; Zou *et al.*, 2021) [3, 16, 19, 20]. We witnessed the consequences of this first-hand, experiencing symptoms such as fatigue, dizziness and heaviness of body. In addition, there were also indirect exposure examples, which involved the consumption of products coming from affected agriculture ecosystems like fruits, vegetables, fish, and seafood (Wang *et al.*, 2017; Zheng *et al.*, 2020) [16, 19].

Cadmium is not very essential minute for life of many things. Besides, the burning of fossil

fuels another pollutant that is not discouraged by nature is also a threat to the environment (Hirao-Suzuki *et al.*, 2021; Karami *et al.*, 2022; Sun *et al.*, 2021; Wang *et al.*, 2017) [4, 7, 13, 16]. The reverses cadmium threat will rape all through the woman pregnancy due to the nature of the toxin being able to diffuse from the placenta and reach the fetus (Ibiwoye *et al.*, 2019) [5]. Also, during the course of gestation this metal can enter and not leave blood, but will head for breast milk - where reproductive organs and blood vessels (Ibiwoye *et al.*, 2019; Liu *et al.*, 2020; Sola *et al.*, 2022) [5, 9, 12] get impacted in babies. Cream focuses on the overall health of a person (liver, kidneys, intestine) and other special organs. (Hirao-Suzuki *et al.*, 2021; Yuan *et al.*, 2021; Zheng *et al.*, 2020) [4, 18, 19]. Intrinsically, cadmium is a cumulative contamination, showing pathogenic effects and the structural modification of the cellular tissues of different cells (Wang *et al.*, 2017) [16]. It is even worse and, additionally, it can attack the skeleton, the tissue of the bone, the stem cells (Wang *et al.*, 2017; Zheng *et al.*, 2020) [16, 19]. The first experiments showed the possibility of cadmium transference to the cells of the reprogrammed testes and especially of the reprogrammed tubule, as well as the risk of cadmium effects on the testes in fashion (Hirao-Suzuki *et al.*, 2021; Liu *et al.*, 2020; Wang *et al.*, 2017) [4, 9, 16]. The impact of nuclear radiation on an individual biological tool the nucleus is the interruption of translation into the DNA and RNA of every exerciser (Dong *et al.*, 2021; Hirao-Suzuki *et al.*, 2021; Suzuki *et al.*, 2017; Yuan *et al.*, 2021) [2, 4, 14, 18]. For this purpose, it can be considered that cadmium is tissue of a disastrous and dangerous solution, and it is carcinogenic.

Leek, the part of the crop that is usually utilized across the world. It has plenty of endowable advantages, in addition to its natural fibers, due to its high content of the active biologically constituents (Venditti *et al.*, 2021) [15]. Thanks to its structure being rich in sulfur compounds, leek could be useful in have antibacterial property (Venditti *et al.*, 2021) [15]. Polyphenols material of the plant vary from triphenolic acid and ananthochlorin to (Bernaert *et al.*, 2013) [1]. Hence, the objective of the study was to evaluate the possibility of utilization of leek powder in unique proportions in rat diet for the obtaining of positive results in cadmium toxifying and pathogenic effects in rats.

Materials and Methods

Preparation of medicinal plants

Leek leaves (*Allium porrum* L.) were collected from local grocery markets in Najaf, Iraq. The fresh fine leek leaves were selected, sorted, washed with D.W three times (Pierart *et al.*, 2018) [11]. Then, the leaves were spread as a thin layer on perforated aluminum foil (5.50 m x 7.50 m) on the laboratory benches and left for 8 h to air dry at 28±2 °C. The dry leek material was processed by crushing and grinding to be mix with the regular feed (Razi Vaccine and Serum Research Institute, Karaj, Iran).

Cadmium source

Cadmium has been used as (CdCl₂) Gew.379.34 E. Merck, Darmstadt, Germany (Hirao-Suzuki *et al.*, 2021; Sola *et al.*, 2022; Wang *et al.*, 2017; Yuan *et al.*, 2021; Zheng *et al.*, 2020) [4, 12, 16, 18, 19]. Groups T (2,3,4) have been Intraperitoneal Injected ((LD) Lethal dose) 0.4 mg /kg body weight, 5 times per week) (Hirao-Suzuki *et al.*, 2021; Sola *et al.*, 2022; Wang *et al.*, 2017; Yuan *et al.*, 2021; Zheng *et al.*, 2020) [4, 12, 16, 18, 19].

Experimental design and data analysis

We used 28 pairs of rats; the Sprague-Dawley male rats were about 150-180 grams each as our model (Razi Vaccine and Serum Research Institute, Karaj, Iran). The rats been divided in 4 groups of 7 and were kept separately in individual cages with dimensions measured from the side of 380mm by 200mm with 590mm (Sun, *et al.*, 2021; Ibiwoye, *et al.*, 2019) [13, 5]. Rats in the lab have dwelt for a week and named housed to permit them to acclimatize to their new environment (Wang *et al.*, 2017) [16]. With respect to this, Wang *et al.* (2017) [16] enlighten that the rats had been kept under care, inside the managed laboratory condition with the temperature of 22±2 °C, the humidity of 30-35% and the 12-hour light/dark cycle. The trial went eight weeks. Water could be consumed by all the groups in the laboratory who wanted them without being limited to a restricted allowance (Wang and co-workers, 2017) [16]. The treatment was to be allocated randomly to the four groups (Treatments). Traditional balance diet were solely infused either at T₁ (treatment) which were all cadmium fed rats and at T₂ that have their leek leaf diet supplied with talcum powder in their normal feed (4 and 8 percent mixture respectively) while treatments T₃ and T₄ were injected with specialists galvanised diet feed from which rats with 20% portion of leek leaf (20 The feed with being deprived of masticating into solids with appearance and consistence similar to the one in usual eating manner as by accomplish this requirement of using 0.6 liters per 1 kg feed, pressing in the molds and serving out for rats. Weigh of rats from various companies has been measured daily, fluctuations are marked in the quantity of food and water feed, and if any exceptions are noted, an investigation is held.

Experimental measurements

Blood tests

Each rat was given five milliliters of blood after being put to sleep (Lee *et al.*, 2017) [8]. The samples were separated using EDTA tubes with serum Sep clot activator. Kavush Laboratory and the National Iranian Standardization Organization tested the blood.

Alkaline Phosphate Determination:

The liver enzyme ALT (alanine aminotransferase) was tested using the Commercial ALAT (GPT) Test base (DGKC) Method.

$$P - \text{Nitrophenyl phosphate} + H_2O \xrightleftharpoons{ATP} \text{Phosphate} + P - \text{Nitrophenol}$$
 (Mohamed Basuony, Ezar Hafez, Ehab Tousson, Ahmed Massoud, Samar Elsomkhraty and Zoology, 2018) [10].

The AST Determination



Aspartate aminotransferase (AST) was determined using the commercial method ASAT (cot) by the IFCC in order to quantify liver enzyme.

$$L - \text{Aspartate} + 2 - \text{Oxoglutarate} \xrightleftharpoons{ASAT} L - \text{Glutamate} + \text{Oxoglutarate}$$

$$\text{Oxoglutarate} + \text{NADH} + \text{H} \xrightleftharpoons{MDH} L - \text{Malate} + \text{NAD}$$

(Mohamed Basuony, Ezar Hafez, Ehab Tousson, Ahmed Massoud, Samar Elsomkhraty and Zoology, 2018) [10].

Alanine Aminotransferase (ALT) Assessment: The ALT (alanine aminotransferase) was tested using the commercial technique ALAT (GPT) by the IFCC to determine the liver enzyme.



Internal organs: kidneys, heart, testes, and liver

Following the experiment, the rats were put under anesthesia and put to death by euthanasia and the weights of their internal organs were then calculated.

Statistical analysis

In this study, the interaction between the different dosages of leek leaf powder was investigated by topical tests utilizing a complete random design (CRD). Least significant difference (LSD) and general statistics (2012) were employed in the statistical analysis of the data reviewed to analyze statistically significant differences between the treatment averages.

Results

Internal organs

The analyzing demonstrated that cubic lead- based rats with cadmium participating to the positive control resulting in destructive effects on the internal organs whose size was

abnormal as indicated by the different weight of kidneys (5.4 g), testes (5.1 g), liver (19.1 g), heart (0.8 g), and spleen (2.2 g) about the kidney weights of the From the outcomes, it was detected that rats from a T₃ or T₄ group (Supplementation with 20% or 30% leek powder, respectively) are resistant to cadmium negative effects. T₃ and T₄ increases in intraabdominal organ weight and led too kidney weights of 3.2 g and 3.6 g, testes weight of 5.4 g and 5.5 g, liver weight of 11.9 g and 12.1 g, heart weight of 1g and 0.9 g and spleen weights of 0.8 g and Table 1. Effect of injection with cadmium and feeding on a diet supplemented with leek (*Allium porrum* L.) powder on rat's Internal organs: 3 replications are things like: Heart, abdominal, Kidney, and Testicles, *Values are means of this. Experiments will be carried out in seven different tanks with the first two tanks (T₁ and T₂) containing animals injected with cadmium and the third (T₃) and fourth (T₄) tank containing cadmium-injected animals fed on a diet supplemented with 2% and 3% leek powder, respectively. The Duncan's Multiple Ranges test shows the "P=0.05" level is being significantly differed by ($p \leq 0.05$) within a column in the case of the same letter or letters. The result of the analysis has found the weight quantities of kidneys, liver and spleen all raise, the evidence show kidney poisoning, inflammatory inside organs (Figure 1).The effect of cadmium on both the weight of the heart and testicles leads to a decrease in their weight, and this is evidence of a decrease in their efficiency and atrophy of these organs.




Fig 1: Shown evidence of a state of inflammation and poisoning at the level of the internal organs of the body.

Table 1: Effect of injection with cadmium and feeding on diet supplemented with leek (*Allium porrum* L.) powder on rat's internal organs: Heart, Liver, Spleen, Testicles, and Kidney

Treatments (T)	Parameters					
	Kidney weight	Testis Weight	Liver Weight	Heart weight	Spleen weight	
T ₁	2.8a±0	7.5a±0	10.0a±0	1.2a±0	0.7a±0	
T ₂	5.4d±0	5.1 c±0	19.1d±0	0.8c±0	2.2c±0	
T ₃	3.2b±0	5.4ab±0	11.9b±0	1.0b±0	0.8a±0	
T ₄	3.6c±0	5.5bc±0	12.1 c±0	0.9bc±0	1.2b±0	
P-Value	Treatment	<.0001	<.0001	<.0001	<.0001	<.0001
	Time (Ti)	<.0001	<.0001	<.0001	<.0001	<.0001
	T*Ti	<.0001	<.0001	<.0001	<.0001	<.0001

*Values are means of 3 replications. Treatments are control (T₁), positive control T₂ (animal injected with cadmium), and T₃ and T₄ for cadmium-injected animals fed on a diet supplemented with 2% and 3% leek powder, respectively. Means followed by the different letter(s) with in a column are significantly different according to Duncan's multiple ranges test ($p \leq 0.05$)

Figure 1. Cadmium negative effect on the rat's internal organs weights. *Values are means of 3 replications. Treatments are control (T₁), positive control T₂ (animal

injected with cadmium), and T₃ and T₄ for cadmium-injected animals fed on a diet supplemented with 2% and 3% leek powder, respectively. Bars that have different letter

(s) are significantly different according to Duncan's multiple ranges test ($p \leq 0.05$).

Blood cadmium levels and liver enzymes (AST, ALT, and Alkaline Phosphatase)

It was observed that the highest levels of cadmium in rat's blood ($25.7 \mu\text{g/L}$) were recorded in T₂ (positive untreated control), while the control T₁ recorded $0.4 \mu\text{g/L}$ which was the lowest among the treatments. On the other hand, leek leaf powder was effective in decreasing cadmium levels in

rat's blood compared to T₂. The therapeutic treatments T₃ and T₄ resulted in cadmium levels of $5.3 \mu\text{g/L}$ and $5.6 \mu\text{g/L}$, respectively. The liver enzymes (AST, ALT, Alkaline Phosphatase) were highly affected by cadmium (Figure 2). Cadmium in the T₂ treatment led to a significant increase in the levels of these enzymes compared to the T₁ control and the groups fed on leek powder T₃ and T₄ treatments. The latter two treatments significantly lowered the enzyme levels (Table 2) under study.

Table 2: Cadmium negative effect on the rat's blood parameters: cadmium [$\mu\text{g/L}$], Alkaline Phosphatase, Alanine Transaminase and Aspartate Transaminase level.

Treatments (T)		Parameters			
		Cadmium [$\mu\text{g/L}$]	AST (U/l)	ALT (U/l)	ALP (IU/L)
T ₁		0.4a±0	22.5a±0	29.1a±0	18.3a±0
T ₂		25.7d±0	60.1d±0	117.0c±0	49.0d±0
T ₃		5.3b±0	29.4b±0	37.2b±0	20.6b±0
T ₄		5.6c±0	33.7c±0	37.9b±0	21.3c±0
P-Value	T	<.0001	<.0001	<.0001	<.0001
	Time	<.0001	<.0001	<.0001	<.0001
	T*Ti	<.0001	<.0001	<.0001	<.0001

*Values are means of 3 replications. Treatments are control (T₁), positive control T₂ (animal injected with cadmium), and T₃ and T₄ for cadmium-injected animals fed on a diet supplemented with 2% and 3% leek powder, respectively. Duncan's multiple ranges test indicates that the means in a column that are followed by a different letter or letters are statistically different ($p \leq 0.05$)




Fig 2: Cadmium negative effect on the rat's blood parameters: cadmium [$\mu\text{g/L}$], Alkaline Phosphatase (Alk. Ph), Alanine Transaminase (ALT) and Aspartate Transaminase (AST) level. Values are means of 3 replications. The animal injected with cadmium (positive control T₂), control (T₁), and cadmium-injected animals provided a meal supplemented with 2% and 3% leek powder, respectively, are the treatments. Duncan's multiple ranges test indicates that bars with distinct letter(s) are significantly different ($p \leq 0.05$)

Discussion

The discussion part could focus on the implications related to the results, showing the comparisons with the existing precedents, giving possible explanations for the observed effects and suggesting possible conclusions about the overall implications. This is the space for discussing your findings and the context of research questions as well as the chances to share others' insights and produce a more scientific view on a topic. Concerning the influence of cadmium on the targeted body organs that came out as results in agreement position with other researches in the field by cadmium (Ibiwoye *et al.*, 2019; Wang *et al.*, 2017) [5, 16]. Your results discussion would detail the importance of this data, look at other works on the topic, give possible

reasons for the findings and talk about the general implication these findings might have. This is the place to translate your findings in line with the primary research question and give as much of your own insight as you might into the whole scientific study of the motive considered. Apart from that, there are also other substantial aspects to talk about such as how these results affect the field in particular and the applications that are possible in the future (Venditti *et al.*, 2021) [15].

Conclusion

The investigation showed the negative effects of cadmium on the weights of internal organs and various liver enzymes, in addition to the levels of cadmium in the blood. The study

found that Leek powder has an effective role in reducing the effects caused by cadmium, the treatment T₄ showed very good results, while the best results were the result of T₃ in reducing those toxic effects of cadmium. These results showed the Leek important role for health by reducing the negative effects heavy metals (Cadmium).

Authors' contributions

The authors collaborated with each other to complete the study. Mortada designed the study and supervised its implementation. Mortada and Ali performed the laboratory analyses, as well as preparing the manuscript first draft. Zinah and Hasan analyzed the data and reviewed the literature for this study. All authors have read the study (final version) and confirmed it for publication

Conflict interest

Neither the authors nor the researchers involved in this study have any financial or personal interests that could potentially lead to conflicts of interest. The research was conducted with a focus solely on contributing to the scientific knowledge and understanding within the field.

Funding

This entire study wasn't funded.

Acknowledgement

The authors thank the University of Kufa.

References

- Bernaert N, De Clercq H, Van Bockstaele E, De Loose M, Van Droogenbroeck B. Antioxidant changes during postharvest processing and storage of leek (*Allium ampeloprasum* var. porrum). *Postharvest Biology and Technology*. Belgium. 2013;86:8-16. DOI: 10.1016/j.postharvbio.2013.06.010.
- Dong W, Liu G, Zhang K, Tan Y, Zou H, Yuan Y, *et al.* Cadmium exposure induces rat proximal tubular cells injury via p62-dependent Nrf2 nucleus translocation mediated activation of AMPK/AKT/mTOR pathway. *Ecotoxicology and Environmental Safety*; c2021. DOI: 10.1016/j.ecoenv.2021.112058.
- Hayat MT, Nauman M, Nazir N, Ali S, Bangash N. Environmental Hazards of Cadmium: Past, Present, and Future. In: *Cadmium Toxicity and Tolerance in Plants: From Physiology to Remediation*. Elsevier Inc.; c2018. DOI: 10.1016/B978-0-12-814864-8.00007-3.
- Hirao-Suzuki M, Takeda S, Sakai G, Waalkes MP, Sugihara N, Takiguchi M, *et al.* Cadmium-stimulated invasion of rat liver cells during malignant transformation: Evidence of the involvement of oxidative stress/TET₁-sensitive machinery. *Toxicology*. Japan; c2021. DOI: 10.1016/j.tox.2020.152631.
- Ibiwoye MO, Matthews Q, Travers K, Foster JD. Association of Acute, High-dose Cadmium Exposure with Alterations in Vascular Endothelial Barrier Antigen Expression and Astrocyte Morphology in the Developing Rat Central Nervous System. *Journal of Comparative Pathology*; c2019. DOI: 10.1016/j.jcpa.2019.08.008.
- Kang J, Jin W, Wang J, Sun Y, Wu X, Liu L, *et al.* Antibacterial and anti-biofilm activities of peppermint essential oil against *Staphylococcus aureus*. *Lwt*. Elsevier. 2019;101(November 2018):639-645. DOI: 10.1016/j.lwt.2018.11.093.
- Karami E, Goodarzi Z, Ghanbari A, Dehdashti A, Bandegi AR, Yosefi S, *et al.* Dataset on biochemical markers and histological alterations in rat kidney intoxicated with cadmium chloride and treated with antioxidant. *Data in Brief*; c2022. DOI: 10.1016/j.dib.2022.108394.
- Lee J, Shin J, Kim J. Effect of Blood Decontamination on Orthodontic Bracket Bonding. *The Journal of the Korean Academy of Pediatric Dentistry*. Dankook, South Korea; c2017. DOI: 10.5933/jkapd.2017.44.3.341.
- Liu J, Zeng L, Zhuang S, Zhang C, Li Y, Zhu J, *et al.* Cadmium exposure during prenatal development causes progesterone disruptors in multiple generations via steroidogenic enzymes in rat ovarian granulosa cells. *Ecotoxicology and Environmental Safety*; c2020. DOI: 10.1016/j.ecoenv.2020.110765.
- Mohamed B, Hafez E, Tousson E, Ahmed M, Samar Elsomkhraty SE, *Zoology*. 2018 Beneficial role of *Panax ginseng* root aqueous.pdf. *American Journal of Biological Chemistry*. Tanta, Egypt; c2018. DOI: <https://www.researchgate.net/publication/274371974> Beneficial.
- Pierart A, Dumat C, Maes AQM, Roux C, Sejalon-Delmas N. Opportunities and risks of biofertilization for leek production in urban areas: Influence on both fungal diversity and human bioaccessibility of inorganic pollutants. *Science of the Total Environment*. France; c2018. DOI: 10.1016/j.scitotenv.2017.12.100.
- Sola E, Moyano P, Flores A, García J, García JM, Anadon MJ, *et al.* Cadmium-induced neurotoxic effects on rat basal forebrain cholinergic system through thyroid hormones disruption. *Environmental Toxicology and Pharmacology*; c2022. DOI: 10.1016/j.etap.2021.103791.
- Sun Y, Zong C, Liu J, Zeng L, Li Q, Liu Z, *et al.* C-myc promotes miR-92a-2-5p transcription in rat ovarian granulosa cells after cadmium exposure. *Toxicology and Applied Pharmacology*; c2021. DOI: 10.1016/j.taap.2021.115536.
- Suzuki M, Takeda S, Teraoka-Nishitani N, Yamagata A, Tanaka T, Sasaki M, *et al.* Cadmium-induced malignant transformation of rat liver cells: Potential key role and regulatory mechanism of altered apolipoprotein E expression in enhanced invasiveness. *Toxicology*; c2017. DOI: 10.1016/j.tox.2017.03.014.
- Venditti M, Ben Rhouma M, Romano MZ, Messaoudi I, Reiter RJ, Minucci S, *et al.* Evidence of melatonin ameliorative effects on the blood-testis barrier and sperm quality alterations induced by cadmium in the rat testis. *Ecotoxicology and Environmental Safety*. Elsevier Inc. 2021;226:112878. DOI: 10.1016/j.ecoenv.2021.112878.
- Wang YJ, Yan J, Zou XL, Guo KJ, Zhao Y, Meng CY, *et al.* Bone marrow mesenchymal stem cells repair cadmium-induced rat testis injury by inhibiting mitochondrial apoptosis. *Chemico-Biological Interactions*; c2017. DOI: 10.1016/j.cbi.2017.04.024.
- Yu Y, Yu L, Zhou X, Qiao N, Qu D, Tian F, *et al.* Effects of acute oral lead exposure on the levels of essential elements of mice: A metallomics and dose-

- dependent study. *Journal of Trace Elements in Medicine and Biology*, 2020, 62(April).
DOI: 10.1016/j.jtemb.2020.126624.
18. Yuan J, Zhao Y, Bai Y, Gu J, Yuan Y, Liu X, *et al.* Cadmium induces endosomal / lysosomal enlargement and blocks autophagy flux in rat hepatocytes by damaging microtubules. *Ecotoxicology and Environmental Safety*. 2021;228:112993.
DOI: 10.1016/j.ecoenv.2021.112993.
19. Zheng J, Zhuo L, Ran D, Ma Y, Luo T, Zhao H, *et al.* Cadmium induces apoptosis via generating reactive oxygen species to activate mitochondrial p53 pathway in primary rat osteoblasts. *Toxicology*; c2020.
DOI: 10.1016/j.tox.2020.152611.
20. Zou H, Wang L, Zhao J, Yuan Y, Wang T, Bian J, *et al.* MiR-155 promotes cadmium-induced autophagy in rat hepatocytes by suppressing Rheb expression. *Ecotoxicology and Environmental Safety*. 2021;227:112895.
DOI: 10.1016/J.ECOENV.2021.112895.